

DOCUMENT RESUME

ED 126 392

CG 010 681

AUTHOR Karpowitz, Dennis H.
TITLE Predicting Deviancy in Family Interaction.
PUB DATE 75
NOTE 8p.; Paper presented at the Annual Meeting of the Rocky Mountain Psychological Association (Salt Lake City, Utah, May 1975)

EDRS PRICE MF-\$0.83 HC-\$1.67 Plus Postage.
DESCRIPTORS Behavioral Science Research; *Behavior Patterns; Family Life; *Family Relationship; *Interaction; *Interpersonal Relationship; *Parent Child Relationship; Research Projects; Speeches; *Stimulus Behavior

ABSTRACT

Should a stimulus be defined as the single event immediately preceding a response (simple stimulus) or as a constellation of antecedents representing several preceding events (complex stimulus)? Sixty-eight families with a child between four and eight years of age were observed, and family interactions coded in the naturalistic setting of the home. A behavioral coding system permitted rapid sequential recording of behaviors. Results were significant in two cross validation groups. The findings were: (1) In all cases the immediately preceding stimulus predicted the type of ensuing response better than single stimuli two or three steps removed. (2) Simple and complex stimuli were both predictive of children's deviant responses. (3) Complex stimuli were often better predictors of children's nondeviant behavior than were simple stimuli. (4) Little more than four percent of the possible complex stimulus combinations accounted for more than 80% of the total stimulus behaviors. (Author)

* Documents acquired by ERIC include many informal unpublished *
* materials not available from other sources. ERIC makes every effort *
* to obtain the best copy available. Nevertheless, items of marginal *
* reproducibility are often encountered and this affects the quality *
* of the microfiche and hardcopy reproductions ERIC makes available *
* via the ERIC Document Reproduction Service (EDRS). EDRS is not *
* responsible for the quality of the original document. Reproductions *
* supplied by EDRS are the best that can be made from the original. *

ED126392

Predicting Deviancy in Family Interaction

Dennis H. Karpowitz

University of Kansas

U.S. DEPARTMENT OF HEALTH,
EDUCATION & WELFARE
NATIONAL INSTITUTE OF
EDUCATION

THIS DOCUMENT HAS BEEN REPRODUCED EXACTLY AS RECEIVED FROM THE PERSON OR ORGANIZATION ORIGINATING IT. POINTS OF VIEW OR OPINIONS STATED DO NOT NECESSARILY REPRESENT OFFICIAL NATIONAL INSTITUTE OF EDUCATION POSITION OR POLICY.

Brief of a Paper Presented at the Rocky Mountain Psychological Association Convention, Salt Lake City, Utah, May, 1975

Abstract

Should a stimulus be defined as the single event immediately preceding a response (simple stimulus) or as a constellation of antecedents representing several preceding events (complex stimulus)? Sixty-eight families with a child between four and eight years of age were observed and family interactions coded in the naturalistic setting of the home. A behavioral coding system permitted rapid sequential recording of behaviors. Results were significant in two cross validation groups.

- a. In all cases the immediately preceding stimulus predicted the type of ensuing response better than single stimuli two or three steps removed.
- b. Simple and complex stimuli were both predictive of children's deviant responses.
- c. Complex stimuli were often better predictors of children's nondeviant behavior than were simple stimuli.
- d. Little more than four percent of the possible complex stimulus combinations accounted for more than 80 percent of the total stimulus behaviors.

Purpose

The purpose of the present social interaction study was to compare the relative utility of a simple stimulus with that of a complex stimulus in predicting the naturally occurring behavior of children. Prior studies of stimulus control have typically limited themselves to studying the function only of the stimulus immediately preceding the relevant response. Yet, it has long been recognized that stimulus constellations or stimuli in sequences may provide greater predictive power than a simple stimulus (e.g., Hull, 1929).

Procedure

A total of 68 families with a child between four and eight years of age were recruited either by advertising or direct invitation. Each family was observed for five days, forty-five minutes per day. A revision of the observational coding system developed by Patterson, Ray, Shaw, and Cobb (1969) was employed. The system was designed for rapid sequential recording of the child's behavior, the responses of family members, the child's response, etc. Spearman-Brown corrected observer agreement coefficients ranged from .91 to .98. The data were sorted into the framework of a sequential process model for stimulus constellations. To improve the probability of finding results which indicated relatively stable proportions, only cells were analyzed which met a minimum frequency requirement. Predictive ability was defined in terms of conditional probabilities (Patterson, 1973). All results were based on cross-subject analyses. The sample was randomly divided into two groups. "t" tests were used to compare differences and only differences which were significant ($p < .05$) in both groups were reported.

Results

Seventy percent of the possible stimulus constellations actually occurred. Only four percent of the constellations met the stability criteria. However, these four percent of the constellations accounted for more than 80% of the recorded behaviors. In all cases the immediately preceding simple stimulus better predicted the type of ensuing response than simple stimuli two or three steps removed from the response. Simple and complex stimuli were both predictive of children's deviant behavior. Complex stimuli were often better predictors of children's non-deviant behavior than were simple stimuli.

Conclusions and Implications

Children's deviant behavior may be cued or "set off" by a simple, single stimulus. On the other hand non-deviant behavior in children often seems to be cued or "set off" by a much more complex chain of events. Programs to modify children's behavior by increasing the frequency of non-deviant behavior may be improved by establishing chains of (positive) cues rather than just a single cue.

References

- Hull, C.L. A functional interpretation of the conditioned reflex. Psychological Review, 1929, 36, 498-511.
- Patterson, G.R., Changes in status of family members as controlling stimuli: a basis for describing treatment process. In L.A. Hammerlynck, L.C. Handy, and E.J. Mash (Eds.), Behavior Change: Methodology, concepts and practice. Champaign, Ill.: Research Press, 1973, 169-191.
- Patterson, G.R., Ray, R.S., Shaw, D.A., & Cobb, J.A. Manual for coding family interactions, sixth revision. Available from ASIS National Auxillary Publication Service, in care of CCM Information Service, Inc., 909 Third Avenue, New York, New York 10022. Document number 01234, 1969.

Table 1

| <u>Agents</u> | | <u>Behaviors</u> | |
|----------------------|---|------------------|--|
| Target Child | | Deviant | |
| | | Nondeviant | |
| Father (F) | } | Positive (+) | |
| Mother (M) | | Neutral (0) | |
| Older Sibling (OS) | | Negative (-) | |
| Younger Sibling (YS) | | | |

Table 2

| | 10" Interval # 1 | 10" Interval # 2 | 10" Interval # 3 |
|--------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------|------------------|
| BEHAVIORAL EVENTS: | Mother pats Johnney on the head → Johnney smiles → Father says, "You are a good boy" → Johnney smiles → Little Sister teases Johnney → Johnney hits little sister | | |

CODED SEQUENCE:

M₁ → TCND₁ → F₂ → TCND₂ → YS-3 → TCD₃

COMPARISONS: 1a.

b.

F₂

YS-3 → TCD₃

c.

M₁

TCD₃

2.

M₁

F₂

YS-

TCD

Table 3

Sequences of Three Antecedents for Which the Proportion
Deviant Behavior was Significantly Different from
the Immediately Preceding Stimulus Alone

| Stimulus Sequence (n-3, n-2, n-1) | N | t score |
|------------------------------------------------------------------------|----|----------|
| Father Positive, Father Positive, Father Positive | 67 | 2.418* |
| Father Neutral, Father Positive, Father Positive | 68 | -2.232* |
| Father Positive, Father Neutral, Father Positive | 65 | -2.048* |
| Father Positive, Father Positive, Father Neutral | 68 | -3.328* |
| Father Neutral, Father Positive, Father Neutral | 67 | -1.998** |
| Father Neutral, Father Neutral, Father Neutral | 68 | 5.376*** |
| Mother Positive, Mother Positive, Mother Positive | 68 | 3.040** |
| Mother Positive, Mother Neutral, Mother Neutral | 53 | 2.040* |
| Mother Positive, Mother Negative, Mother Negative | 39 | 3.322** |
| Father Positive, Older Sibling Positive, Older Sibling Positive | 29 | 3.739*** |
| Older Sibling Neutral, Older Sibling Neutral, Older Sibling Neutral | 21 | 3.287** |
| Father Neutral, Father Neutral Younger Sibling Positive | 41 | 2.659* |
| Father Neutral, Younger Sibling Positive, Younger Sibling Positive | 40 | 2.145* |

* = $p < .05$ ** = $p < .01$ *** = $p < .001$